



Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV.

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 8/31/2020

Visit the AIDSinfo website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <https://aidsinfo.nih.gov/e-news>.

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 8/31/2020

Appendix A. List of Abbreviations (Last updated May 7, 2013; last reviewed May 7, 2013)

Acronym/Abbreviation	Definition
ABGs	arterial blood gasses
ACTG	AIDS Clinical Trials Group
AFB	acid-fast bacilli
AIN	anal intraepithelial neoplasia
ALT	alanine aminotransferase
anti-HBc	hepatitis B core antibody
anti-HBs	hepatitis B surface antibody
ART	antiretroviral therapy
ARV	antiretroviral
ASCCP	American Society for Colposcopy and Cervical Pathology
ASC-H	atypical squamous cells—cannot exclude high grade cervical squamous intraepithelial lesion
ASC-US	atypical squamous cells of uncertain significance
AST	serum aspartate aminotransferase
AUC	area under the curve
BA	bacillary angiomatosis
BAL	bronchoalveolar lavage
BID	twice a day
BIW	twice a week
CAP	community-acquired pneumonia
CAPD	continuous ambulatory peritoneal dialysis
CD4	CD4 T lymphocyte cell
CDC	Centers for Disease Control and Prevention
CDI	<i>Clostridium difficile</i> -associated infection
CES-D	Center for Epidemiologic Studies Depression Scale
CFU	colony-forming unit
CIA	chemiluminescence immunoassays
CIN	cervical intraepithelial neoplasia
C_{\max}	maximum concentration
C_{\min}	minimum concentration
CMV	cytomegalovirus
CNS	central nervous system
CPE	central nervous system penetration effectiveness
CrCl	creatinine clearance
CSF	cerebrospinal fluid
CT	computed tomography

CYP3A4	Cytochrome P450 3A4
DAAs	direct acting antiviral agents
DOT	directly observed therapy
DS	double strength
EDTA	ethylenediaminetetraacetic acid
EIAs	enzyme immunoassays
EM	erythema multiforme
FDA	Food and Drug Administration
FTA-ABS	fluorescent treponemal antibody absorbed
g	gram
G6PD	Glucose-6-phosphate dehydrogenase
GFR	glomerular filtration rate
GI	gastrointestinal
HAV	hepatitis A virus
HBV	hepatitis B virus
HCV	hepatitis C virus
HHV-8	human herpesvirus-8
HPA	hypothalamic-pituitary-adrenal
HPV	human papillomavirus
HSIL	high grade cervical squamous intraepithelial lesion
HSV	herpes simplex virus
HSV-1	herpes simplex virus 1
HSV-2	herpes simplex virus 2
ICP	intracranial pressure
ICU	intensive care unit
IFN	interferon
IgG	immunoglobulin G
IgM	immunoglobulin M
IGRA	interferon-gamma release assays
IM	intramuscular
IND	investigational new drug
IRIS	immune reconstitution inflammatory syndrome
IRU	immune recovery uveitis
IV	intravenous
IVIG	intravenous immunoglobulin
JCV	JC virus
KS	Kaposi Sarcoma
LEEP	loop electrosurgical excision procedure
LP	lumbar puncture
LSIL	low grade squamous intraepithelial lesion

LTBI	latent tuberculosis infection
MAC	<i>Mycobacterium avium</i> complex
MAI	<i>Mycobacterium avium intracellulare</i>
MCD	multicentric Castleman's disease
MDR TB	multi-drug-resistant tuberculosis
mg	milligram
mmHg	millimeters of mercury
MSM	men who have sex with men
MTB	<i>Mycobacterium tuberculosis</i>
NAA	nucleic acid amplification
NNRTI	non-nucleoside reverse transcriptase inhibitor
NRTI	nucleoside reverse transcriptase inhibitors
NSAID	non-steroidal anti-inflammatory drugs
NVP	nevirapine
OI	opportunistic infection
PCP	<i>Pneumocystis pneumonia</i>
PCR	polymerase chain reaction
PEL	primary effusion lymphoma
PK	pharmacokinetic
PML	progressive multifocal leukoencephalopathy
PO	orally
PORN	Progressive Outer Retinal Necrosis
PPV	polysaccharide vaccine
PSI	pneumonia severity index
q(n)h	every "n" hours
qAM	every morning
QID	four times a day
qPM	every evening
RPR	rapid plasma reagent
RVR	rapid virological response
SCr	serum creatinine
SJS	Stevens-Johnson syndrome
SLE	systemic lupus erythematosus
SQ	subcutaneous
SS	single strength
STD	sexually transmitted disease
SVR	sustained virologic response
TB	tuberculosis
TDM	therapeutic drug monitoring
TE	<i>Toxoplasma</i> encephalitis

TEN	toxic epidermal necrolysis
TID	three times daily
TIW	three times weekly
TP-PA	<i>T. pallidum</i> particle agglutination
TST	tuberculin skin test
ULN	upper limit of normal
VAIN	vaginal intra-epithelial neoplasia
VDRL	Venereal Disease Research Laboratory
VIII	vestibulocochlear
VIN	vulvar intraepithelial neoplasia
VZV	varicella zoster virus
WBC	white blood cell
WHO	World Health Organization
XDR TB	extensively drug-resistant tuberculosis

Abbreviation	Drug Name
3TC	lamivudine
5-FU	fluorouracil
ATV/r	ritonavir-boosted atazanavir
BCA	bichloroacetic acid
BOC	boceprevir
COBI	cobicistat
ddA-TP	dideoxyadenosine triphosphate
ddI	didanosine
DHA	dihydroartemisinin
EFV	efavirenz
EMB	ethambutol
EVG	elvitegravir
FTC	emtricitabine
INH	isoniazid
MVC	maraviroc
PCV13	13-valent pneumococcal conjugate vaccine
PegIFN	peginterferon alfa
PI	protease inhibitor
PPV23	23-valent pneumococcal polysaccharides vaccine
PZA	pyrazinamide
RAL	raltegravir
RBV	ribavirin
RFB	rifabutin
RIF	rifampin

RPT	rifapentine
SMX	sulfamethoxazole
TCA	trichloroacetic acid
TDF	tenofovir disoproxil fumarate
TMP	trimethoprim
TMP-SMX	trimethoprim-sulfamethoxazole
TVR	telaprevir
ZDV	zidovudine